

LaTeX

The plasma-solid transition: two simple examples

V. Celebonovic¹ and W. Dappen²

¹Institute of Physics, Pregrevica 118, 11080 Zemun-Beograd, Yugoslavia

²Dept. of Physics and Astronomy, USC, Los Angeles CA 90089-1342, USA

1. Introduction

The notion of "astrophysical matter" is meant by most people to refer to one of the various forms of plasmas. However, this is not entirely true, because contemporary astrophysics is full of cases of objects built with dominantly high-pressure, solid-state matter. Obvious examples of such dense objects are planetary and stellar interiors [1]. Logically then, between these two regimes, there must exist a region in which a plasma-solid (PS) transition occurs.

The aim of this contribution is to explore in some detail the conditions for the PS transition, using a simple criterion as proposed in [2]. We apply it to the two extreme cases: Fermi-Dirac (FD) and Bose-Einstein (BE) gas. One might object here that the applicability of any of these model systems to real astrophysical objects is not very probable. This is true, but the idea behind the present calculation is to obtain an order-of-magnitude estimate, first in the two pure limits (FD and BE), and later in a mixed FD and BE case.

2. Calculations

The criterion for the existence of the solid phase, proposed in [2], has the form:

$$T \cdot 10^4 T_6^2 \leq \rho \leq 3 \cdot 10^{11} Z^{10} \quad (1)$$

and

$$50 \cdot Z^{-5} T_6^3 \leq \rho \leq 7 \cdot 10^4 T_6^2 \quad (2)$$

where $T_6 = T/10^6$ and ρ is the mass density. Eq.(1) is applicable to "low temperatures".

2.1 The FD case

The equation of state (EOS) of a FD gas has the form:

$$n = \frac{4\pi}{h^3} (2mT)^{3/2} F_{1/2} \left(\frac{\mu}{T} \right) \quad (3)$$

where $k_B = 1$, μ is the chemical potential, and $F_{1/2}$ is the FD integral of the order 1/2 [3].

According to [3]

$$F_{1/2} \left(\frac{\mu}{T} \right) \simeq \frac{2}{3} \mu_0^3 \left[1 + \frac{\pi^4}{48} \left(\frac{T}{\mu_0} \right)^4 \right] \quad (4)$$

where $\mu_0 = Cn^{2/3}$ and $C = (3\pi^2)^{2/3} \frac{\hbar^2}{2m}$. Inserting eq.(4) into eq.(3), it follows that

$$n \simeq \left(\frac{\pi^4}{48} \right)^{3/8} \left(\frac{T}{C} \right)^{3/2} \left(\frac{T^{3/2}}{8} \right)^{3/8} \left(\frac{T^{3/2}}{8 \left(1 - \frac{T^{3/2}}{8} \right)} \right)^{3/8} \quad (5)$$

The mass density ρ and the electron number density n are related by $n = N_A \frac{Z\rho}{A}$. Applying this relation and eq.(5) to eq.(1), one finally gets

$$10^{-8} T^3 \leq \frac{A}{N_A Z} \frac{1}{864^{1/8} 4} \left(\frac{\pi}{C} \right)^{3/2} \left(T^{33/16} + \frac{3}{64} T^{57/16} \right) \leq 3 \cdot 10^{11} Z^{10} \quad (6)$$

which is the allowed interval for the existence of the solid phase of a FD gas.

2.2 The BE case

The EOS of an ideal BE gas has the form [4]

$$n = \frac{g}{\hbar^3} \left(\frac{mT}{2\pi} \right)^{3/2} Li_{3/2} \left[\exp \left(\frac{\mu}{T} \right) \right] \quad (7)$$

where $g = 2s + 1$ is the spin degeneracy factor and $Li_n[z] = \sum_{k=1}^{\infty} \frac{z^k}{k^n}$ is the definition of the polylogarithm. One of the important points of the phase diagram of a BE gas is the Bose-Einstein condensation (BEC) point, defined as the zero point of the chemical potential.

Inserting this condition into eq.(7),and performing the necessary algebra,one gets the following result for the BEC temperature

$$T_{BEC} = \frac{2\pi\hbar^2}{[g\zeta(3/2)]^{2/3}} \frac{n^{2/3}}{m} \quad (8)$$

(n - the number density of a BE gas ; ζ - the Riemann zeta function).Solving eq.(8) for the number density and inserting the result into eq.(1), it follows that

$$\frac{2\pi\hbar^2}{(10^{16}m^5)^{1/3}} \frac{T^2}{[g\zeta(3/2)]^{2/3}} \leq T_{BEC} \leq 2\pi\hbar^2 \left(\frac{3 \cdot 10^{11}}{g\zeta(3/2)} \right)^{2/3} \frac{1}{m^{5/3}} \quad (9)$$

This inequality gives the interval of temperatures in which the BEC temperature has to be in order for the BEC to occur in a solid phase.In deriving it,we have used $\rho = mn$.

3. Discussion and conclusions

In this contribution we have discussed the plasma-solid transition for two model systems: the FD and the BE gas.The calculations were performed according to a criterion proposed by Kirzhnits,and the results are given in eqs.(6) and (9).These two expressions are to be considered as limiting cases of astrophysical reality.Of course,in the most common astrophysical systems,such as stellar interiors and atmospheres,one encounters mixture of the general form BE and FD.The problem of a PS transition in such systems will be treated elsewhere.

However,our eq.(9) is applicable in the exotic albeit real astrophysical setting of white dwarfs and neutron stars (and similar other dense objects).There are indications that a PS transition might occur in them [5].Comparing the results of eq.(9) with the temperature profiles of neutron stars,in the domain after the PS transition,one could determine the depth at which BEC could possibly occur in these objects.This is just a qualitative indication of the possible applicability of eq.(9),and the details will be considered at length elsewhere.

References

- [1] E.Schatzman and F.Praderie: *Astrophysique:Les Etoiles*, InterEditions/Editions du CNRS,Paris (1990).
- [2] D.A.Kirzhnits: *Zh.Exp.Theor.Fiz.*,**38**,503 (1960).
- [3] V.Celebonovic: *Publ.Astron.Obs.Belgrade*,**60**,16 (1998).
- [4] L.E.Reichl: *A Modern Course in Statistical Physics*,Edward Arnold (Publishers) Ltd.,London (1988).
- [5] D.G.Yakovlev,K.P.Levenfish and Yu.A.Shibanov: *Cooling Neutron stars and Superfluidity in Their Interiors*,preprint astro-ph/9906456 (1999).